

## Adult Resuscitation Trolley Arrest Drugs Checklist

### Drugs Drawer:

#### Loose / Individual Cardiac Arrest Drugs as below.

*Pre-filled syringe delivery is preference to mini-jets or ampoules where possible.*

|   |   |
|---|---|
| <b>Anaphylaxis box</b> (sealed by pharmacy)   | <b>One</b>                                |
| <b>Adrenaline pre-filled syringe 1 mg in 10 ml (1:10,000)</b>   | <b>Six</b>                                |
| <b>Amiodarone pre-filled syringe 300 mg in 10 ml</b>  | <b>Two</b>                                |
| <b>Atropine pre-filled syringe – at least 3 mg available</b><br>(Potential options depending on pharmacy availability)<br>(or 3mg in 10 ml – if 1 mg not available from Pharmacy)   | <b>One</b>                                |
| <b>Calcium Chloride pre-filled syringe 10% in 10 ml (preferred choice)</b><br><u>alternatively</u><br><b>Calcium Gluconate 10% 30mls</b>  | <b>One</b>                                |
| <b>Sodium Bicarbonate 8.4%</b><br>(dependant on pharmacy/supply availability);<br><br>- <b>polyfusor 200 ml x 1 (preferred choice)</b><br><u>alternatively</u><br>- <b>ampoules (IV injection) 10 mls (come in box of 10)</b> | <b>One</b>                                |
| <b>Magnesium Sulphate 50% ampoule 5g in 10 ml (2 mmol/ml)</b><br>(supplied as box of 10)  | <b>One</b>                                |
| <b>Naloxone</b><br><br>- <b>pre-filled syringe 400 mcg in 1 ml</b><br><u>or</u><br>- <b>ampoules 400 mcg</b>  | <b>Five minimum</b><br>– (total 2000 mcg) |
| <b>In IV fluid drawer:</b>  |   |
| <b>Glucose IV 10% 500 ml bag</b>  | <b>One</b>                                |

PLEASE TURN OVER

The clinical areas below require **X2 500ml bags of Intralipid 20%** (reversal for local anaesthetic toxicity) in the IV fluid drawer;

- Theatres/PACU
- SDAU/Nettlewell Assessment Unit
- Labour Ward
- Alexandra Day Stay Unit (ADSU)
- Penn Ward
- Tye Green Ward
- ITU/HDU
- Harold Ward
- Henry Moore
- Kingsmoor Ward
- Bevan Ward

All of the following areas will have or receive patients with spinal/epidural procedures and therefore require **Intralipid 20%** to be available for emergency use.

### Reminder

**Please also make sure you have checked your HYPOGLYCAEMIA EMERGENCY BOX. This is NOT to be kept on the Resuscitation Trolley.**